

University of Groningen

A translational approach to the genetics of renal disease

Doorenbos, Carolina Roelfiena Cornelia

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Doorenbos, C. R. C. (2012). *A translational approach to the genetics of renal disease*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 3

Possible renoprotection by vitamin D in chronic renal disease: beyond mineral metabolism

CRC Doorenbos, J van den Born, GJ Navis, MH de Borst

Nature Reviews Nephrology. 2009 Dec;5(12):691-700.

Abstract

Vitamin D is typically viewed as a key player in the regulation of calcium and phosphate levels and the control of bone metabolism; however, growing evidence suggests that vitamin D deficiency may also have an important role in the progressive loss of renal function. Vitamin D deficiency is particularly frequent in patients with chronic kidney disease, in whom it is associated with increased mortality. Studies indicate that treatment with vitamin D analogues reduces proteinuria, suppresses the renin-angiotensin-aldosterone system (RAAS), and exerts anti-inflammatory and immunomodulatory effects. These pleiotropic effects render vitamin D a potentially interesting treatment modality for renoprotection in patients with chronic kidney disease. Whether vitamin D has clinically relevant renoprotective effects in addition to RAAS blockade is currently under investigation.

Introduction

Vitamin D has been recognized for decades as a key player in the control of bone metabolism through the regulation of calcium and phosphate homeostasis. This hormone regulates serum concentrations of calcium and phosphate by stimulating the absorption of these minerals in the intestine and their reabsorption by the kidney. Vitamin D deficiency results in increased circulating levels of parathyroid hormone (PTH), which has several disadvantageous effects on bone metabolism. PTH activates osteoblasts to stimulate the maturation of osteoclasts. Mature osteoclasts degrade the mineralized collagen matrix in bone, resulting in osteopenia, osteoporosis, and increased risk of fracture. Another important consequence of high circulating levels of PTH levels is phosphaturia, which can result in low levels of serum phosphate. Consequently, the calcium-phosphorus product decreases, which results in decreased mineralization of the collagen matrix and can lead to the development of rickets in children and osteomalacia in adults (1).

The decline in the incidence of rickets that was observed following the fortification of food products with vitamin D indicated that the key problems associated with vitamin D deficiency were solved. Over the past decade, however, two important findings have questioned this assumption. First, despite fortification of foods with vitamin D in Western countries, a considerable proportion of the population remained insufficient or even deficient in vitamin D. Certain subpopulations such as elderly individuals and patients with chronic kidney disease (CKD) are at particular risk of vitamin D deficiency. Second is the increasing awareness that vitamin D not only has a role in the regulation of bone metabolism, but also affects several other organ systems. For example, kidneys express functional vitamin D receptors (VDRs) and respond to treatment with active vitamin D. A growing number of experimental studies suggest the vitamin D axis has a renoprotective role. Maintenance of adequate serum vitamin D levels might, therefore, be of particular importance for patients with renal disease, not only for the prevention of bone disease, but also for the protection of renal function.

This Review first outlines the physiology of vitamin D and discusses the physiological processes that might be affected in patients with CKD. The epidemiology of vitamin D deficiency as well as its implications in both the general population and in patients with renal disease is also described. The main focus is on the pathogenic consequence of vitamin D deficiency in the kidney and on the renoprotective potential of pharmacological strategies that use vitamin D analogues.

Vitamin D physiology

In the presence of solar ultraviolet B radiation, vitamin D₃ is photochemically produced in the skin from 7-dehydrocholesterol (Figure 3.1). Alternatively, vitamin D₃ can be absorbed through the intestine from food sources such as oily fish, fortified dairy products and mushrooms. Ingested or photosynthesized vitamin D₃ is converted to active vitamin D (1 α ,25(OH)₂ vitamin D₃ or calcitriol, subsequently referred to in this Review as active vitamin D), via two hydroxylation steps. First, vitamin D₃ is converted by 25-hydroxylase in the liver to form 25(OH) vitamin D₃ (calcidiol). Calcidiol is subsequently converted to active vitamin D in the proximal tubular epithelial cells of the kidney. Transport of the different vitamin D metabolites to the liver, kidney, and target organs is facilitated by

the plasma vitamin D-binding protein (DBP) (2). The calcidiol-DBP complex is filtered by the glomerulus before it enters the proximal tubule cells via megalin-mediated uptake at the brush border (2). In tubular epithelial cells, the conversion of calcidiol into active vitamin D is regulated by 25(OH) vitamin D₃ 1 α -hydroxylase (CYP27B1). This enzyme has an important role in the regulation of circulating concentrations of active vitamin D, and is controlled by several peptide hormones and regulatory feedback loops. PTH is the main inducer of 1 α -hydroxylase activity. PTH is produced by calcium-sensing cells of the parathyroid gland, mainly under hypocalcemic conditions. In addition, 1 α -hydroxylase activity is induced by calcitonin (3), a hormone that is produced by the C cells of the thyroid gland under normocalcemic conditions when the PTH system is downregulated.

Circulating levels of active vitamin D are negatively regulated by several feedback mechanisms (Figure 3.1). First, vitamin D can be directly inactivated by the enzyme 1,25-dihydroxyvitamin D₃ 24-hydroxylase (CYP24A1) by the addition of another hydroxyl (OH) group at position 24. This step results in the generation of 1,24,25(OH)₃ vitamin D₃ and ultimately produces the water-soluble calcitroic acid (1 α -hydroxy-23-carboxy-24,25,-26,27-tetranorvitamin D₃), which is excreted by the kidney. Second, 1 α -hydroxylase expression is downregulated by active vitamin D itself via vitamin D-responsive elements in the promoter region of 1 α -hydroxylase (4). Third, active vitamin D stimulates the production of fibroblast growth factor (FGF)23 in osteocytes (5). This growth factor acts via the FGF receptor 1 (FGFR-1)-Klotho complex, which functions as an FGF23 receptor (6) to negatively regulate renal synthesis of 1 α -hydroxylase (Figure 3.1) (7).

Several factors also influence circulating levels of active vitamin D. The effects of seasonal changes (8) and sunlight exposure (9) on levels of active vitamin D have been recognized for decades. Circulating vitamin D levels also differ with ethnicity, being lowest in non-Hispanic blacks (10). Diet is also a contributing factor, but the contribution of diet to overall levels of active vitamin D is limited because much higher amounts are produced in the skin (1).

The actions of active vitamin D are mediated by the VDR. This receptor is part of the nuclear receptor superfamily and is predominantly located in the nucleus, although in some cells it can be associated with caveolae in the plasma membrane. Following the recruitment of transcriptional coactivators, the activated VDR binds to vitamin D-response elements in the promoter region of target genes. An overview of the diverse VDR target genes that have been identified in several cell types has been published elsewhere (9–15).

The VDR also exerts nongenomic actions. For example, the VDR can form a complex with the p65 subunit of NF κ B to produce anti-inflammatory actions (16). Another nongenomic mechanism of action of VDR is achieved by so-called rapid responses. This mechanism has been reviewed elsewhere (17). Briefly, binding of active vitamin D to caveolae-associated VDRs activates second messenger systems, which may force the opening of voltage-gated calcium or chloride channels, and facilitate cross-talk with the nucleus to modulate the expression of target genes. The VDR is expressed by more than 30 organ systems and tissues (18–20), including the kidney. Available data on the expression pattern of VDRs in the normal kidney are limited, but immunohistochemically detectable levels are expressed by proximal and distal tubular epithelial cells, glomerular parietal epithelial cells, and collecting duct cells (21). Treatment of cultured tubular epithelial cells (16), mesangial cells (22), podocytes (23), and juxtaglomerular cells (23), with ac-




Figure 3.1: Vitamin D physiology. Vitamin D₃ is generated from 7-dehydrocholesterol in the skin under the influence of UV-B radiation or is obtained from dietary sources by absorption through the intestine. Circulating vitamin D₃ is transported in conjugation with DBP. In the liver, vitamin D₃ is converted to 25(OH) vitamin D₃ by 25-hydroxylase. 25(OH) vitamin D₃ is then converted to the active vitamin D metabolite, 1α,25(OH)₂ vitamin D₃ in the kidney by the enzyme 1α-hydroxylase. 1α-hydroxylase is regulated by several mechanisms, including PTH, calcitonin, and FGF23. Active vitamin D can be deactivated by addition of another hydroxyl (OH) group at the position 24, after which the end product is excreted. Abbreviations: 24-OHase, 1,25-dihydroxyvitamin D₃ 24-hydroxylase; DBP, vitamin D-binding protein; FGF, fibroblast growth factor; PTH, parathyroid hormone; UV, ultraviolet; vit, vitamin. Permission obtained from Nature Publishing Group Ltd © Deeb, K. K. *et al.* Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat. Rev. Cancer* 7, 684-700 (2007).

tive vitamin D or vitamin D analogues such as paricalcitol, modulated the expression of target genes, suggesting the presence of a functional VDR in these cells. Whether renal VDR expression is modulated in patients with CKD, is unknown.

Epidemiology of vitamin D deficiency

The definition of vitamin D deficiency is still a matter of debate. Some consider serum levels of 25(OH) vitamin D₃ < 75 nmol/l (< 30 ng/ml) to be insufficient, levels < 37 nmol/l (< 15 ng/ml) deficient, and levels < 25 nmol/l (< 10 ng/ml) severely deficient (24). However, most experts define vitamin D deficiency as 25(OH) vitamin D₃ levels < 50 nmol/l (< 20 ng/ml) (1, 25–27). A review of several reports of vitamin D intoxication suggests that serum levels of 25(OH) vitamin D₃ > 375–500 nmol/l (> 150–200 ng/ml) are associated with a risk of hypercalcemia and should be considered toxic (1, 28).

Data from consecutive cohort studies of the general population from the National Health and Nutrition Examination Survey (NHANES) suggested that changes had occurred in levels of serum vitamin D in the US population between the periods of 1988–1994 (NHANES III) and 2001–2004 (10). This change might be related to the increased use of sunscreen (29), decreased amounts of outdoor activity (30), and the increased prevalence of obesity (31), all of which are associated with reduced vitamin D levels. Although these data raise considerable concerns, the validity of comparing different measurements of 25(OH) vitamin D₃ has been questioned because of methodological issues that can arise due to calibration problems, lack of standard operating procedures and the use of a variety of equipment (32–34). Adjusting for methodological changes in the measurement of vitamin D levels demonstrated that much of the observed difference in vitamin D levels between the NHANES III and NHANES 2001–2004 studies could be accounted for by changes unrelated to actual changes in vitamin D status (35). Until a reference measurement procedure for 25(OH) vitamin D₃ is available, caution must be taken with the interpretation of epidemiological data on vitamin D (36).

Patients with CKD, in particular those on dialysis, generally have more severe vitamin D deficiency than the general population (10, 37–39). Vitamin D insufficiency (defined as levels < 75 nmol/l [< 30 ng/ml]) and severe vitamin D deficiency (defined as levels < 25 nmol/l [< 10 ng/ml]) are both more common in patients with predialysis CKD than in the normal population (Figure 3.2) (37). Some, but not all, studies have found a positive association between stage of CKD and prevalence of vitamin D deficiency (40, 41). Predialysis CKD patients who live in a sunny country are less likely to be vitamin D insufficient or deficient than those who live in less sunny areas (42). In a cross-sectional cohort of 825 dialysis patients in the US, only 22% had 25(OH) vitamin D₃ levels of > 75 nmol/l (> 30 ng/ml), while 60% had levels between 25 and 75 nmol/l (10 and 30 ng/ml) (43). In a French study of dialysis patients, 42% had severe vitamin D deficiency (Figure 3.2) (38). An analysis of patients on peritoneal dialysis showed similar percentages (44). Thus, in patients with end-stage renal disease, vitamin D deficiency is an even more prevalent and severe problem than it is in patients with mild to moderate renal insufficiency.




Figure 3.2: Prevalence of vitamin D deficiency in the general population and in patients with chronic kidney disease. Prevalence of vitamin D levels > 75 nmol/l (> 30 ng/ml) (defined by some researchers as sufficient), between 25 and 75 nmol/l (10 and 30 ng/ml), and < 25 nmol/l (< 10 ng/ml) (severely deficient) in the a — general population (8), b,c — in patients with preterminal chronic kidney disease (stages 3 and 4) (37), and d — hemodialysis patients (stage 5) (38). Vitamin D insufficiency is common in all categories of patients, whereas severe deficiency seems to be more prevalent in those with more severe renal disease. Abbreviations: CKD, chronic kidney disease; GFR, glomerular filtration rate.

Implications of vitamin D deficiency

In the general population, vitamin D deficiency has been independently associated with a small but significantly increased risk of all-cause mortality (45–47). Other studies found that vitamin D deficiency may be associated with cardiovascular morbidity (48) and mortality (49). A subgroup analysis showed that the strongest associations between vitamin D deficiency and mortality occurred in patients without cardiovascular disease, hypertension, or diabetes. This finding may imply that vitamin D deficiency exerts detrimental actions before cardiovascular disease is established. Whether vitamin D treatment could provide cardiovascular protection has been the subject of intense debate since the publication of negative findings by Hsia *et al.* (50) in 2007. Of note, researchers have commented that the vitamin D replacement dose used in that study may have been inadequate to achieve an effect (51).

The relationship between vitamin D deficiency and impaired bone metabolism is illustrated by the role of vitamin D deficiency in the development of rickets and osteomalacia (52). Several studies have documented beneficial effects of vitamin D supplementation (≥ 700 –800 IU daily in combination with calcium) in patients with osteoporosis (53–56). Lower doses of vitamin D₃, or doses that do not lead to the restoration of adequate serum 25(OH) vitamin D₃ levels, may not effectively lower fracture risk (25, 57).

In the NHANES III cohort, vitamin D deficiency was associated with an increased risk of albuminuria (58). Whether vitamin D deficiency is the cause or consequence (through reduced reuptake of the calcidiol-DBP complex in tubular cells under proteinuric conditions, as is discussed below) of albuminuria, or whether it is an epiphenomenon, is the subject of ongoing randomized, controlled trials (59). In patients with CKD, several studies have demonstrated an independent inverse association between serum levels of active vitamin D and the rate of renal function loss or mortality (41, 60, 61). Similarly, Wolf *et al.* reported an association between vitamin D deficiency and early mortality in a cohort of patients on dialysis (43). Importantly, subsequent treatment of these patients with active vitamin D was associated with increased survival.

Low serum vitamin D levels are more strongly associated with increased risk of coronary atherosclerosis in patients with CKD than in individuals with normal renal function (62). In a population of patients on peritoneal dialysis, the risk of cardiovascular morbidity was higher in individuals with vitamin D deficiency than in those without (44). In other studies, vitamin D deficiency was independently associated with several cardiovascular risk factors such as hypertension, insulin resistance, diabetes, and dyslipidemia (63, 64). These associations could at least in part explain the increased incidence of cardiovascular morbidity and mortality in patients with CKD and vitamin D deficiency. Osteoporosis and osteopenia are common in patients with CKD (65, 66). This detrimental effect on bone is most likely related to disturbed vitamin D and PTH metabolism in these patients.

The prevalence of vitamin D deficiency in patients with CKD and the association of vitamin D deficiency with increased mortality, cardiovascular morbidity and bone disease, warrants adequate correction of vitamin D levels well before these patients develop end-stage renal disease. The Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines advise vitamin D supplementation in patients with stages 3–4 CKD if serum 25(OH) vitamin D₃ level is < 75 nmol/l (< 30 ng/ml) to reduce the risk of low bone mineral density and hip fracture (67). Increasing understanding of the potential renoprotective

effects of active vitamin D might provide an additional reason for the early initiation of vitamin D supplementation in patients with renal disease. Emerging data suggests that correction of vitamin D may potentially slow the progression of renal disease in these patients.

Vitamin D deficiency in renal disease

Mechanisms of reduced vitamin D levels

After filtration by the glomerulus, the reuptake of 25(OH) vitamin D₃ (bound to DBP), takes place at the brush border of the proximal tubular epithelial cells (Figure 3.3). This process is mediated by megalin, and can result in release of either active 1,25(OH)₂ vitamin D₃ or inactive 25(OH) vitamin D₃ into the circulation on the basolateral side. Although reduced renal megalin expression has not been demonstrated in patients with renal disease, studies in cultured tubular epithelial cells suggest that chronic exposure to albumin reduces megalin expression (68). In addition, renal megalin messenger RNA (mRNA) expression is reduced in models of experimental nephrotic syndrome *in vivo* (69). Furthermore, megalin knockout mice develop low-molecular-weight proteinuria and actually lose large amounts of vitamin D-DBP in their urine (70). These findings are supported by studies in a kidney-specific megalin knockout mouse-when fed a vitamin D-depleted diet, these mice demonstrated lower circulating levels of 25(OH) vitamin D₃ and strongly elevated 25(OH) vitamin D₃-DBP excretion than did wildtype mice (71). Furthermore, kidney-specific megalin knockout mice were hypocalcemic and had osteomalacia (softening of the bones) (71). In a rat model of nephrotic syndrome induced by puromycin aminonucleoside, serum levels of 25(OH) vitamin D₃ decreased concomitantly with increased proteinuria (72). Since liver mRNA expression of both CYP2R1 (the protein product of which metabolizes vitamin D₃ to 25(OH) vitamin D₃) and the gene that encodes DBP were unchanged, the decreased levels of 25(OH) vitamin D₃ might not be a result of dysfunctional hydroxylation or transport to the liver. Rather, these studies indicate that 25(OH) vitamin D₃-DBP might be lost into the urine in the nephrotic state. Alternatively, the reduced serum levels of 25(OH) vitamin D₃ in CKD patients might occur secondary to decreased photoproduction of pre-vitamin D₃ from 7-dehydrocholesterol in the skin because of uremia (73).

Reduced activity of 1 α -hydroxylase is widely accepted to be the main cause of reduced circulating levels of 1,25(OH)₂ vitamin D₃ in patients with CKD (74). In a model of experimental nephrosis, renal mRNA expression of 1-hydroxylase is decreased and CYP24A1 is increased 3 days after induction of nephrosis, resulting in a reduction in serum levels of 1,25(OH)₂ vitamin D₃ (72). Therefore, in patients with CKD, vitamin D deficiency probably results from multiple factors including urinary loss of 25(OH) vitamin D₃-DBP associated with proteinuria (72), reduced activity of 1 α -hydroxylase, and compromised endogenous pre-vitamin D₃ production in the skin (73).

Renoprotective potential of vitamin D

Growing evidence indicates that vitamin D analogues may have beneficial effects in patients with CKD. Antifibrotic effects of vitamin D analogues have been demonstrated in




Figure 3.3: Possible mechanisms of vitamin D deficiency in proteinuric renal disease. a — In healthy individuals, $25(\text{OH})$ vitamin D_3 bound to DBP is filtered through the glomerulus (1). In proximal tubular cells, uptake of $25(\text{OH})$ vitamin $\text{D}_3\text{-DBP}$ is facilitated by megalin. After uptake into lysosomes (2), $25(\text{OH})$ vitamin $\text{D}_3\text{-DBP}$ is either returned to the circulation (3), or converted to active vitamin D by hydroxylation by 1-hydroxylase (4), and released into the circulation (5). b — In patients with proteinuria, megalin is occupied by an extensive protein load that consists mainly of albumin (1); thus, fewer receptors are available to reabsorb $25(\text{OH})$ vitamin $\text{D}_3\text{-DBP}$. In addition, as a consequence of proteinuria, proximal tubular cells are damaged and express less megalin (2), which also reduces $25(\text{OH})$ vitamin $\text{D}_3\text{-DBP}$ reabsorption. 1-hydroxylase expression is also reduced in patients with chronic kidney disease (3), which results in decreased formation of active vitamin D. Abbreviation: DBP, vitamin D-binding protein; vit, vitamin.




Figure 3.4: Overview of the potential renoprotective effects of active vitamin D in the damaged kidney. In the glomerulus, vitamin D protects podocytes by increasing the expression of nephrin and podocin. Under nephrotic conditions, vitamin D reduces the expression of desmin, a marker of podocyte injury. In juxtaglomerular apparatus cells, renin production is inhibited by vitamin D. In macrophages, vitamin D reduces TNF and NFκB expression. Similarly, vitamin D has anti-inflammatory effects in injured tubular epithelial cells, by reducing NFκB availability. Consequently, expression of chemokines such as CCL5 (RANTES), are reduced. Vitamin D can also modify T-helper-cell responses and affect the maturation and differentiation of dendritic cells, which provides protection against inflammatory damage in response to renal injury. Abbreviations: CCL5, CC-chemokine ligand 5; NFκB, nuclear factor κB; T_H, T-helper; TNF, tumor necrosis factor.

cultured renal cells (75) and in animal models of renal damage (76). Several mechanisms could be involved in this renoprotective effect, including direct antiproteinuric effects through the protection of podocytes, interactions with the renin-angiotensin-aldosterone system (RAAS), and anti-inflammatory effects (Figure 3.4).

Antiproteinuric effects

In experimental puromycin aminonucleoside-induced nephrotic syndrome, treatment with active vitamin D or the vitamin D analogue 22-oxacalcitriol has demonstrated clear antiproteinuric effects (72, 77). Administration of active vitamin D or 22-oxacalcitriol partially restored glomerular expression of nephrin, an important structural slit diaphragm protein that is inversely correlated with proteinuria. In podocytes, active vitamin D induces nephrin gene expression (possibly in cooperation with the retinoic acid receptor (78)), probably by binding of the activated VDR to vitamin D response elements in the nephrin gene (79). In addition, vitamin D treatment restores expression of podocin, a molecule associated with the slit diaphragm, and decreases expression of the podocyte injury marker desmin (72). Together, these findings indicate that active vitamin D has a direct protective effect on podocytes, and possibly explains the antiproteinuric effects of vitamin D that have been reported by other studies (80, 81). As proteinuria is a marker of progressive renal dysfunction, these data support the potential of vitamin D administration for long-term renoprotection. Studies to assess the antiproteinuric effects of vitamin D analogues in humans are currently underway. As RAAS blockade is first method of choice for reducing proteinuria, the effects of vitamin D administration together with RAAS blockade are also under investigation (59).

RAAS-related effects

The antiproteinuric effects of vitamin D and its analogues could, at least in part, be explained by actions on the RAAS. Two key publications demonstrated that active vitamin D binds to the promoter region of the renin gene to downregulate renal mRNA expression of renin (68, 82). In VDR knockout mice, inhibition of renin transcription by vitamin D is reduced, which results in RAAS activation and renal injury (23, 82). VDR knockout mice are hypertensive, probably as a consequence of RAAS activation (82). Whether this finding implies that administration of vitamin D analogues could have antihypertensive effects is currently a subject of controversy (83). In cultured juxtaglomerular-like cells, the administration of active vitamin D reduces renin expression by 90% (82) by blocking the cyclic adenosine monophosphate response element in the renin gene promoter (68). In adipocytes, 1,25(OH)₂ vitamin D₃ dose-dependently decreases expression of the gene that encodes the type I angiotensin II receptor (84). Whether VDR and type I angiotensin II receptor interact in renal cells is unknown.

RAAS blockade is currently first-line renoprotection therapy for patients with CKD, but can be associated with a reactive rise in renin (85, 86). Both angiotensin-converting-enzyme (ACE) inhibitors and type I angiotensin II antagonists require concomitant adherence to a low sodium diet and administration of a diuretic to achieve optimal effects on blood pressure and proteinuria (87, 88); however, the benefits of these approaches come at the expense of a potentiated reactive rise in levels of both renin and aldosterone. Data

support direct profibrotic effects of aldosterone (89) and renin by direct effects on the (pro)-renin receptor (90). Thus, the eventual therapeutic benefit of volume depletion by RAAS blockade can be limited by the reactive rise in renin levels. Data from studies in rats support the presence of profibrotic renal effects associated with aggressive volume depletion under RAAS blockade (91). These data, therefore, suggest that blocking the reactive rise in renin during RAAS blockade could have therapeutic benefits. In our opinion, vitamin D is a potential candidate for this purpose. The addition of vitamin D to RAAS blockade therapy might improve the efficacy of RAAS blockade by blunting the reactive rise in renin (and the downstream rise in aldosterone). Indeed, treatment with a combination of paricalcitol and enalapril reduced renal damage in the 5/6 nephrectomy rat model more strongly than did paricalcitol or enalapril alone (92).

Direct inhibition of renin has long been known to have renoprotective potential (93) and may suppress the reactive rise in renin activity. Studies over the past few years have investigated the renoprotective effects achieved by combined treatment with conventional RAAS blockers (that is, ACE inhibitors and angiotensin II receptor blockers) and the direct renin inhibitor aliskiren (94–96). In patients with type 2 diabetes, hypertension, and diabetic nephropathy, combination treatment with aliskiren and losartan for 6 months reduced proteinuria to a greater extent than did losartan alone (94). The reduction in proteinuria seemed to occur independently of blood pressure, which was not significantly affected by treatment. However, the long-term renoprotective effects of combined renin and ACE inhibition or angiotensin II receptor blockade remain to be documented. For the purpose of the current Review, it should be mentioned that direct renin inhibitors affect the enzymatic activity of renin but not its production or its interaction with the prorenin receptor (97). Therefore, inhibition of renin production by vitamin D may still have therapeutic potential for patients with CKD (98).

The anti-inflammatory properties of vitamin D might also add to the value of vitamin D as an adjunct to RAAS blockade, as inflammation is a determinant of RAAS blockade resistance (99, 100). Vitamin D therapy could be particularly beneficial in patients for whom additional RAAS blockade therapy is contraindicated, such as those with hypotension or hyperkalemia.

Anti-inflammatory and immunomodulatory effects

Several studies have demonstrated the anti-inflammatory properties of active vitamin D (16, 22, 101). The concept of vitamin D as an immunomodulatory molecule originated over 25 years ago (102, 103), but the mechanisms involved, and the relevance of these effects to the treatment of renal disease, have only now begun to be understood (104). A study by Zehnder *et al.* reported that serum 1,25(OH)₂ vitamin D₃ levels were inversely associated with renal inflammation in several types of kidney disease (105). As mentioned previously, treatment of cultured proximal tubular epithelial cells with the vitamin D analogue paricalcitol induced the formation of a complex that consisted of VDR and the NF κ B component p65 (16). The formation of this complex led to the reduced binding of NF κ B to the promoter regions of genes involved in the inflammatory response, resulting in decreased activation of genes such as CCL5 (also known as RANTES). Similar results were achieved in mesangial cells-administration of active vitamin D inhibited the activity of CCL2 (also known as MCP-1) by stabilization of the NF κ B inhibitory unit I κ B α (22).

In macrophages, active vitamin D also suppressed NF κ B activity, leading to reduced production of TNF (106). These findings emphasize that VDR regulates the expression of target genes both directly, and also by altering the activity of transcription factors such as NF κ B.

Of interest, in addition to processing in the liver and the kidneys, vitamin D can also be metabolized by cells of the immune system (107). Specifically, activated T cells can convert 25(OH) vitamin D₃ to active vitamin D (107). Moreover, macrophages and some dendritic cells express enzymes required to convert vitamin D₃ into active vitamin D (107–109). The net effect of active vitamin D on T cells is to block the production of cytokines, in particular interferon γ , by T-helper-1 cells, and to promote T-helper-2 cell responses (110). Vitamin D also has inhibitory effects on dendritic cell maturation and differentiation *in vitro*, which results in a marked T-cell hyporesponsiveness (that is, reduced T-cell proliferation and interferon γ production) (111). Vitamin D may have an important role in the maintenance of peripheral tolerance by expanding the pool of antigen-specific regulatory T cells (112, 113). Together, these findings justify studies that investigate the potential of vitamin D analogues for the treatment of inflammatory kidney diseases such as autoimmune nephritis, and for tolerance induction in renal transplant recipients. Preclinical studies in rats have shown improved graft survival following treatment with active vitamin D after kidney transplantation (114). Although cholecalciferol treatment (100,000 IU every 2 weeks for 2 months) in vitamin D-deficient kidney transplant patients has been shown to be safe (115), the dose required to induce renal allograft tolerance or to prolong allograft or patient survival is not known.

Vitamin D could also have important immunomodulatory effects in individuals with non-autoimmune chronic renal disease and in those who do not yet require kidney transplantation. Of note, a recent study demonstrated the importance of renal dendritic cells in the progression of glomerulonephritis after glomerular injury (116). Vitamin D-mediated inhibition of dendritic cell maturation and differentiation might reduce the inflammatory response associated with glomerular injury, which might slow or potentially even halt progression of renal damage.

Conclusions

The kidney has a key role in the metabolism of vitamin D; consequently, renal disease affects vitamin D metabolism. Epidemiological studies indicate that the prevalence of vitamin D deficiency in the general population is considerable, but is higher in patients with CKD. This latter finding is important since low circulating levels of vitamin D have been associated with increased morbidity and mortality, particularly in patients with CKD; however, whether a causal relationship exists remains to be determined.

In addition to the classic effects of vitamin D on calcium and phosphate homeostasis, several studies have revealed other important functions of this hormone. For example, administration of vitamin D in experimental CKD reduces proteinuria, which may relate to the effects of vitamin D on the RAAS and anti-inflammatory actions. Vitamin D decreases RAAS activity by downregulation of renin gene transcription in juxtaglomerular cells. The inhibitory effect on renin is of particular interest, given the reactive rise in renin during RAAS blockade, the cornerstone of current pharmacological treatment in patients

with CKD. Vitamin D might, therefore, be a suitable add-on strategy to RAAS blockade in these patients. Long-term, randomized, controlled trials should address the renoprotective potential of this approach.

The relationship between vitamin D and the immune system is currently under intense investigation. The role of vitamin D in macrophages, T cells and dendritic cells is of particular interest, since these cells have a substantial role in the pathophysiology of renal damage. Results from clinical trials that have used vitamin D analogues in patients with CKD (59) and renal transplant recipients (117) are expected in the near future.

In conclusion, emerging data indicate that vitamin D deficiency has greater clinical consequences than the induction of osteomalacia in non-renal and renal populations. Although the prevalence of rickets has decreased in Western countries, other disease entities that are related to vitamin D deficiency have been recognized. Clinical trials to investigate the role of vitamin D supplementation for renal and cardiovascular protection are currently underway, and will enable the role of vitamin D supplementation in the management of CKD to be better defined.

Acknowledgments

M. H. de Borst is supported by research grants from the Dutch Kidney Foundation (KJPB.08.07) and the University Medical Center Groningen (Mandema stipend).

References

- [1] M. F. Holick. Vitamin d deficiency. *The New England journal of medicine*, 357(3):266–281, Jul 19 2007.
- [2] A. Nykjaer, D. Dragun, D. Walther, H. Vorum, C. Jacobsen, J. Herz, F. Melsen, E. I. Christensen, and T. E. Willnow. An endocytic pathway essential for renal uptake and activation of the steroid 25-(oh) vitamin d3. *Cell*, 96(4):507–515, Feb 19 1999.
- [3] Y. Zhong, H. J. Armbrrecht, and S. Christakos. Calcitonin, a regulator of the 25-hydroxyvitamin d3 1alpha-hydroxylase gene. *The Journal of biological chemistry*, 284(17):11059–11069, Apr 24 2009.
- [4] M. M. Turunen, T. W. Dunlop, C. Carlberg, and S. Vaisanen. Selective use of multiple vitamin d response elements underlies the 1 alpha,25-dihydroxyvitamin d3-mediated negative regulation of the human cyp27b1 gene. *Nucleic acids research*, 35(8):2734–2747, 2007.
- [5] S. Liu, W. Tang, J. Zhou, J. R. Stubbs, Q. Luo, M. Pi, and L. D. Quarles. Fibroblast growth factor 23 is a counter-regulatory phosphaturic hormone for vitamin d. *Journal of the American Society of Nephrology : JASN*, 17(5):1305–1315, May 2006.
- [6] I. Urakawa, Y. Yamazaki, T. Shimada, K. Iijima, H. Hasegawa, K. Okawa, T. Fujita, S. Fukumoto, and T. Yamashita. Klotho converts canonical fgf receptor into a specific receptor for fgf23. *Nature*, 444(7120):770–774, Dec 7 2006.
- [7] T. Shimada, M. Kakitani, Y. Yamazaki, H. Hasegawa, Y. Takeuchi, T. Fujita, S. Fukumoto, K. Tomizuka, and T. Yamashita. Targeted ablation of fgf23 demonstrates an essential physiological role of fgf23 in phosphate and vitamin d metabolism. *The Journal of clinical investigation*, 113(4):561–568, Feb 2004.
- [8] C. Carlberg and S. Seuter. A genomic perspective on vitamin d signaling. *Anticancer Research*, 29(9):3485–3493, Sep 2009.
- [9] S. Christakos, M. Raval-Pandya, R. P. Wernyj, and W. Yang. Genomic mechanisms involved in the pleiotropic actions of 1,25-dihydroxyvitamin d3. *The Biochemical journal*, 316 (Pt 2)(Pt 2): 361–371, Jun 1 1996.
- [10] A. A. Ginde, M. C. Liu, and C. A. Camargo Jr. Demographic differences and trends of vitamin d insufficiency in the us population, 1988-2004. *Archives of Internal Medicine*, 169(6):626–632, Mar 23 2009.
- [11] J. H. White. Profiling 1,25-dihydroxyvitamin d3-regulated gene expression by microarray analysis. *The Journal of steroid biochemistry and molecular biology*, 89-90(1-5):239–244, May 2004.
- [12] Y. Bosse, K. Maghni, and T. J. Hudson. 1alpha,25-dihydroxy-vitamin d3 stimulation of bronchial smooth muscle cells induces autocrine, contractility, and remodeling processes. *Physiological genomics*, 29(2):161–168, Apr 24 2007.
- [13] B. C. Ellis, S. Gattoni-Celli, A. Mancina, and M. S. Kindy. The vitamin d3 transcriptomic response in skin cells derived from the atlantic bottlenose dolphin. *Developmental and comparative immunology*, 33(8):901–912, Aug 2009.
- [14] K. Townsend, V. Trevino, F. Falciani, P. M. Stewart, M. Hewison, and M. J. Campbell. Identification of vdr-responsive gene signatures in breast cancer cells. *Oncology*, 71(1-2):111–123, 2006.
- [15] J. R. Wu-Wong, M. Nakane, J. Ma, X. Ruan, and P. E. Kroeger. Vdr-mediated gene expression patterns in resting human coronary artery smooth muscle cells. *Journal of cellular biochemistry*, 100(6):1395–1405, Apr 15 2007.
- [16] X. Tan, X. Wen, and Y. Liu. Paricalcitol inhibits renal inflammation by promoting vitamin d receptor-mediated sequestration of nf-kappab signaling. *Journal of the American Society of Nephrology : JASN*, 19(9):1741–1752, Sep 2008.

- [17] A. W. Norman. Minireview: vitamin d receptor: new assignments for an already busy receptor. *Endocrinology*, 147(12):5542–5548, Dec 2006.
- [18] A. S. Dusso, A. J. Brown, and E. Slatopolsky. Vitamin d. *American journal of physiology. Renal physiology*, 289(1):F8–28, Jul 2005.
- [19] M. F. Holick and M. Garabedian. *Vitamin D: photobiology, metabolism, mechanism of action, and clinical applications.*, pages 129–137. Primer on the metabolic bone diseases and disorders of mineral metabolism. American Society for Bone and Mineral Research, Washington, DC, 6 edition, 2006. ISBN 9780977888108.
- [20] M. F. Holick. Resurrection of vitamin d deficiency and rickets. *The Journal of clinical investigation*, 116(8):2062–2072, Aug 2006.
- [21] R. Kumar, J. Schaefer, J. P. Grande, and P. C. Roche. Immunolocalization of calcitriol receptor, 24-hydroxylase cytochrome p-450, and calbindin d28k in human kidney. *The American Journal of Physiology*, 266(3 Pt 2):F477–85, Mar 1994.
- [22] Z. Zhang, W. Yuan, L. Sun, F. L. Szeto, K. E. Wong, X. Li, J. Kong, and Y. C. Li. 1,25-dihydroxyvitamin d3 targeting of nf-kappab suppresses high glucose-induced mcp-1 expression in mesangial cells. *Kidney international*, 72(2):193–201, Jul 2007.
- [23] Z. Zhang, L. Sun, Y. Wang, G. Ning, A. W. Minto, J. Kong, R. J. Quigg, and Y. C. Li. Renoprotective role of the vitamin d receptor in diabetic nephropathy. *Kidney international*, 73(2):163–171, Jan 2008.
- [24] R. Vieth. The role of vitamin d in the prevention of osteoporosis. *Annals of Medicine*, 37(4): 278–285, 2005.
- [25] H. A. Bischoff-Ferrari, E. Giovannucci, W. C. Willett, T. Dietrich, and B. Dawson-Hughes. Estimation of optimal serum concentrations of 25-hydroxyvitamin d for multiple health outcomes. *The American Journal of Clinical Nutrition*, 84(1):18–28, Jul 2006.
- [26] P. Lips. Which circulating level of 25-hydroxyvitamin d is appropriate? *The Journal of steroid biochemistry and molecular biology*, 89-90(1-5):611–614, May 2004.
- [27] A. Malabanan, I. E. Veronikis, and M. F. Holick. Redefining vitamin d insufficiency. *Lancet*, 351(9105):805–806, Mar 14 1998.
- [28] G. Jones. Pharmacokinetics of vitamin d toxicity. *The American Journal of Clinical Nutrition*, 88(2):582S–586S, Aug 2008.
- [29] L. Y. Matsuoka, L. Ide, J. Wortsman, J. A. MacLaughlin, and M. F. Holick. Sunscreens suppress cutaneous vitamin d3 synthesis. *The Journal of clinical endocrinology and metabolism*, 64(6): 1165–1168, Jun 1987.
- [30] R. Scragg and C. A. Camargo Jr. Frequency of leisure-time physical activity and serum 25-hydroxyvitamin d levels in the us population: results from the third national health and nutrition examination survey. *American Journal of Epidemiology*, 168(6):577–86; discussion 587–91, Sep 15 2008.
- [31] J. Wortsman, L. Y. Matsuoka, T. C. Chen, Z. Lu, and M. F. Holick. Decreased bioavailability of vitamin d in obesity. *The American Journal of Clinical Nutrition*, 72(3):690–693, Sep 2000.
- [32] N. Binkley, D. Krueger, C. S. Cowgill, L. Plum, E. Lake, K. E. Hansen, H. F. DeLuca, and M. K. Drezner. Assay variation confounds the diagnosis of hypovitaminosis d: a call for standardization. *The Journal of clinical endocrinology and metabolism*, 89(7):3152–3157, Jul 2004.
- [33] R. J. Singh. Are clinical laboratories prepared for accurate testing of 25-hydroxy vitamin d? *Clinical chemistry*, 54(1):221–223, Jan 2008.

- [34] A. M. Yates, A. Bowron, L. Calton, J. Heynes, H. Field, S. Rainbow, and B. Keevil. Interlaboratory variation in 25-hydroxyvitamin d2 and 25-hydroxyvitamin d3 is significantly improved if common calibration material is used. *Clinical chemistry*, 54(12):2082–2084, Dec 2008.
- [35] A. C. Looker, C. M. Pfeiffer, D. A. Lacher, R. L. Schleicher, M. F. Picciano, and E. A. Yetley. Serum 25-hydroxyvitamin d status of the us population: 1988-1994 compared with 2000-2004. *The American Journal of Clinical Nutrition*, 88(6):1519–1527, Dec 2008.
- [36] G. D. Carter. 25-hydroxyvitamin d assays: the quest for accuracy. *Clinical chemistry*, 55(7): 1300–1302, Jul 2009.
- [37] R. E. LaClair, R. N. Hellman, S. L. Karp, M. Kraus, S. Ofner, Q. Li, K. L. Graves, and S. M. Moe. Prevalence of calcidiol deficiency in ckd: a cross-sectional study across latitudes in the united states. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 45(6):1026–1033, Jun 2005.
- [38] G. Jean, B. Charra, and C. Chazot. Vitamin d deficiency and associated factors in hemodialysis patients. *Journal of renal nutrition : the official journal of the Council on Renal Nutrition of the National Kidney Foundation*, 18(5):395–399, Sep 2008.
- [39] E. A. Gonzalez, A. Sachdeva, D. A. Oliver, and K. J. Martin. Vitamin d insufficiency and deficiency in chronic kidney disease. a single center observational study. *American Journal of Nephrology*, 24 (5):503–510, Sep-Oct 2004.
- [40] A. Levin, G. L. Bakris, M. Molitch, M. Smulders, J. Tian, L. A. Williams, and D. L. Andress. Prevalence of abnormal serum vitamin d, pth, calcium, and phosphorus in patients with chronic kidney disease: results of the study to evaluate early kidney disease. *Kidney international*, 71(1): 31–38, Jan 2007.
- [41] D. V. Barreto, F. C. Barreto, S. Liabeuf, M. Temmar, F. Boitte, G. Choukroun, A. Fournier, and Z. A. Massy. Vitamin d affects survival independently of vascular calcification in chronic kidney disease. *Clinical journal of the American Society of Nephrology : CJASN*, 4(6):1128–1135, Jun 2009.
- [42] L. Cuppari, A. B. Carvalho, and S. A. Draibe. Vitamin d status of chronic kidney disease patients living in a sunny country. *Journal of renal nutrition : the official journal of the Council on Renal Nutrition of the National Kidney Foundation*, 18(5):408–414, Sep 2008.
- [43] M. Wolf, A. Shah, O. Gutierrez, E. Ankers, M. Monroy, H. Tamez, D. Steele, Y. Chang, C. A. Camargo Jr, M. Tonelli, and R. Thadhani. Vitamin d levels and early mortality among incident hemodialysis patients. *Kidney international*, 72(8):1004–1013, Oct 2007.
- [44] A. Y. Wang, C. W. Lam, J. E. Sanderson, M. Wang, I. H. Chan, S. F. Lui, M. M. Sea, and J. Woo. Serum 25-hydroxyvitamin d status and cardiovascular outcomes in chronic peritoneal dialysis patients: a 3-y prospective cohort study. *The American Journal of Clinical Nutrition*, 87 (6):1631–1638, Jun 2008.
- [45] A. Zittermann, S. Iodice, S. Pilz, W. B. Grant, V. Bagnardi, and S. Gandini. Vitamin d deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *The American Journal of Clinical Nutrition*, 95(1):91–100, Jan 2012.
- [46] P. Autier and S. Gandini. Vitamin d supplementation and total mortality: a meta-analysis of randomized controlled trials. *Archives of Internal Medicine*, 167(16):1730–1737, Sep 10 2007.
- [47] M. L. Melamed, B. Astor, E. D. Michos, T. H. Hostetter, N. R. Powe, and P. Muntner. 25-hydroxyvitamin d levels, race, and the progression of kidney disease. *Journal of the American Society of Nephrology : JASN*, 20(12):2631–2639, Dec 2009.
- [48] T. J. Wang, M. J. Pencina, S. L. Booth, P. F. Jacques, E. Ingelsson, K. Lanier, E. J. Benjamin, R. B. D’Agostino, M. Wolf, and R. S. Vasan. Vitamin d deficiency and risk of cardiovascular disease. *Circulation*, 117(4):503–511, Jan 29 2008.

- [49] H. Dobnig, S. Pilz, H. Scharnagl, W. Renner, U. Seelhorst, B. Wellnitz, J. Kinkeldei, B. O. Boehm, G. Weihrauch, and W. Maerz. Independent association of low serum 25-hydroxyvitamin d and 1,25-dihydroxyvitamin d levels with all-cause and cardiovascular mortality. *Archives of Internal Medicine*, 168(12):1340–1349, Jun 23 2008.
- [50] J. Hsia, G. Heiss, H. Ren, M. Allison, N. C. Dolan, P. Greenland, S. R. Heckbert, K. C. Johnson, J. E. Manson, S. Sidney, M. Trevisan, and Women’s Health Initiative Investigators. Calcium/vitamin d supplementation and cardiovascular events. *Circulation*, 115(7):846–854, Feb 20 2007.
- [51] D. E. Wallis and S. Penckofer. Letter by wallis and penckofer regarding article, ”calcium/vitamin d supplementation and cardiovascular events”. *Circulation*, 116(3):e86; author reply e87, Jul 17 2007.
- [52] J. M. Pettifor. *Vitamin D deficiency and nutritional rickets in children.*, volume 1 of *Vitamin D*, pages 1065–1084. Elsevier Academic Press, Burlington, MA, 2, illustrated edition, 2005. ISBN 9780122526893.
- [53] A. Avenell, W. J. Gillespie, L. D. Gillespie, and D. O’Connell. Vitamin d and vitamin d analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane database of systematic reviews (Online)*, (2)(2):CD000227, Apr 15 2009.
- [54] B. M. Tang, G. D. Eslick, C. Nowson, C. Smith, and A. Bensoussan. Use of calcium or calcium in combination with vitamin d supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet*, 370(9588):657–666, Aug 25 2007.
- [55] M. C. Chapuy, M. E. Arlot, F. Duboeuf, J. Brun, B. Crouzet, S. Arnaud, P. D. Delmas, and P. J. Meunier. Vitamin d3 and calcium to prevent hip fractures in the elderly women. *The New England journal of medicine*, 327(23):1637–1642, Dec 3 1992.
- [56] B. Dawson-Hughes, S. S. Harris, E. A. Krall, and G. E. Dallal. Effect of calcium and vitamin d supplementation on bone density in men and women 65 years of age or older. *The New England journal of medicine*, 337(10):670–676, Sep 4 1997.
- [57] A. M. Grant, A. Avenell, M. K. Campbell, A. M. McDonald, G. S. MacLennan, G. C. McPherson, F. H. Anderson, C. Cooper, R. M. Francis, C. Donaldson, W. J. Gillespie, C. M. Robinson, D. J. Torgerson, W. A. Wallace, and RECORD Trial Group. Oral vitamin d3 and calcium for secondary prevention of low-trauma fractures in elderly people (randomised evaluation of calcium or vitamin d, record): a randomised placebo-controlled trial. *Lancet*, 365(9471):1621–1628, May 7-13 2005.
- [58] I. H. de Boer, G. N. Ioannou, B. Kestenbaum, J. D. Brunzell, and N. S. Weiss. 25-hydroxyvitamin d levels and albuminuria in the third national health and nutrition examination survey (nhanes iii). *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 50(1):69–77, Jul 2007.
- [59] H. J. Lambers Heerspink, R. Agarwal, D. W. Coyne, H. H. Parving, E. Ritz, G. Remuzzi, P. Audhya, M. J. Amdahl, D. L. Andress, and D. de Zeeuw. The selective vitamin d receptor activator for albuminuria lowering (vital) study: study design and baseline characteristics. *American Journal of Nephrology*, 30(3):280–286, 2009.
- [60] D. Inaguma, H. Nagaya, K. Hara, M. Tatematsu, H. Shinjo, S. Suzuki, T. Mishima, and K. Kurata. Relationship between serum 1,25-dihydroxyvitamin d and mortality in patients with pre-dialysis chronic kidney disease. *Clinical and experimental nephrology*, 12(2):126–131, Apr 2008.
- [61] P. Ravani, F. Malberti, G. Tripepi, P. Pecchini, S. Cutrupi, P. Pizzini, F. Mallamaci, and C. Zoccali. Vitamin d levels and patient outcome in chronic kidney disease. *Kidney international*, 75(1):88–95, Jan 2009.
- [62] I. H. de Boer, B. Kestenbaum, A. B. Shoben, E. D. Michos, M. J. Sarnak, and D. S. Siscovick. 25-hydroxyvitamin d levels inversely associate with risk for developing coronary artery calcification. *Journal of the American Society of Nephrology : JASN*, 20(8):1805–1812, Aug 2009.

- [63] M. Chonchol and R. Scragg. 25-hydroxyvitamin d, insulin resistance, and kidney function in the third national health and nutrition examination survey. *Kidney international*, 71(2):134–139, Jan 2007.
- [64] D. Martins, M. Wolf, D. Pan, A. Zadshir, N. Tareen, R. Thadhani, A. Felsenfeld, B. Levine, R. Mehrotra, and K. Norris. Prevalence of cardiovascular risk factors and the serum levels of 25-hydroxyvitamin d in the united states: data from the third national health and nutrition examination survey. *Archives of Internal Medicine*, 167(11):1159–1165, Jun 11 2007.
- [65] R. Marcen, C. Caballero, O. Uriol, A. Fernandez, J. J. Villafruela, J. Pascual, J. Martins, N. Rodriguez, F. J. Burgos, and J. Ortuno. Prevalence of osteoporosis, osteopenia, and vertebral fractures in long-term renal transplant recipients. *Transplantation proceedings*, 39(7):2256–2258, Sep 2007.
- [66] V. Pichette, A. Bonnardeaux, L. Prudhomme, M. Gagne, J. Cardinal, and D. Ouimet. Long-term bone loss in kidney transplant recipients: a cross-sectional and longitudinal study. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*, 28(1):105–114, Jul 1996.
- [67] Kidney disease outcomes quality initiative clinical practice guidelines for bone metabolism and disease in chronic kidney disease [online], 2003.
- [68] W. Yuan, W. Pan, J. Kong, W. Zheng, F. L. Szeto, K. E. Wong, R. Cohen, A. Klopot, Z. Zhang, and Y. C. Li. 1,25-dihydroxyvitamin d3 suppresses renin gene transcription by blocking the activity of the cyclic amp response element in the renin gene promoter. *The Journal of biological chemistry*, 282(41):29821–29830, Oct 12 2007.
- [69] X. Tan, Y. Li, and Y. Liu. Paricalcitol attenuates renal interstitial fibrosis in obstructive nephropathy. *Journal of the American Society of Nephrology : JASN*, 17(12):3382–3393, Dec 2006.
- [70] J. R. Leheste, B. Rolinski, H. Vorum, J. Hilpert, A. Nykjaer, C. Jacobsen, P. Aucouturier, J. O. Moskaug, A. Otto, E. I. Christensen, and T. E. Willnow. Megalin knockout mice as an animal model of low molecular weight proteinuria. *The American journal of pathology*, 155(4):1361–1370, Oct 1999.
- [71] J. R. Leheste, F. Melsen, M. Wellner, P. Jansen, U. Schlichting, I. Renner-Muller, T. T. Andreassen, E. Wolf, S. Bachmann, A. Nykjaer, and T. E. Willnow. Hypocalcemia and osteopathy in mice with kidney-specific megalin gene defect. *The FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, 17(2):247–249, Feb 2003.
- [72] I. Matsui, T. Hamano, K. Tomida, K. Inoue, Y. Takabatake, Y. Nagasawa, N. Kawada, T. Ito, H. Kawachi, H. Rakugi, E. Imai, and Y. Isaka. Active vitamin d and its analogue, 22-oxacalcitriol, ameliorate puromycin aminonucleoside-induced nephrosis in rats. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*, 24(8):2354–2361, Aug 2009.
- [73] A. I. Jacob, A. Sallman, Z. Santiz, and B. W. Hollis. Defective photoproduction of cholecalciferol in normal and uremic humans. *The Journal of nutrition*, 114(7):1313–1319, Jul 1984.
- [74] D. L. Andress. Vitamin d in chronic kidney disease: a systemic role for selective vitamin d receptor activation. *Kidney international*, 69(1):33–43, Jan 2006.
- [75] Y. Li, B. C. Spataro, J. Yang, C. Dai, and Y. Liu. 1,25-dihydroxyvitamin d inhibits renal interstitial myofibroblast activation by inducing hepatocyte growth factor expression. *Kidney international*, 68(4):1500–1510, Oct 2005.
- [76] U. Schwarz, K. Amann, S. R. Orth, A. Simonaviciene, S. Wessels, and E. Ritz. Effect of 1,25 (oh)₂ vitamin d3 on glomerulosclerosis in subtotaly nephrectomized rats. *Kidney international*, 53(6):1696–1705, Jun 1998.

- [77] H. Xiao, W. Shi, S. Liu, W. Wang, B. Zhang, Y. Zhang, L. Xu, X. Liang, and Y. Liang. 1,25-dihydroxyvitamin d(3) prevents puromycin aminonucleoside-induced apoptosis of glomerular podocytes by activating the phosphatidylinositol 3-kinase/akt-signaling pathway. *American Journal of Nephrology*, 30(1):34–43, 2009.
- [78] M. Okamura, Y. Takano, Y. Saito, J. Yao, and M. Kitamura. Induction of nephrin gene expression by selective cooperation of the retinoic acid receptor and the vitamin d receptor. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*, 24(10):3006–3012, Oct 2009.
- [79] K. Yamauchi, Y. Takano, A. Kasai, K. Hayakawa, N. Hiramatsu, N. Enomoto, J. Yao, and M. Kitamura. Screening and identification of substances that regulate nephrin gene expression using engineered reporter podocytes. *Kidney international*, 70(5):892–900, Sep 2006.
- [80] Y. Wang, J. Zhou, A. W. Minto, B. K. Hack, J. J. Alexander, M. Haas, Y. C. Li, C. W. Heilig, and R. J. Quigg. Altered vitamin d metabolism in type ii diabetic mouse glomeruli may provide protection from diabetic nephropathy. *Kidney international*, 70(5):882–891, Sep 2006.
- [81] A. Kuhlmann, C. S. Haas, M. L. Gross, U. Reulbach, M. Holzinger, U. Schwarz, E. Ritz, and K. Amann. 1,25-dihydroxyvitamin d3 decreases podocyte loss and podocyte hypertrophy in the subtotaly nephrectomized rat. *American journal of physiology.Renal physiology*, 286(3):F526–33, Mar 2004.
- [82] Y. C. Li, J. Kong, M. Wei, Z. F. Chen, S. Q. Liu, and L. P. Cao. 1,25-dihydroxyvitamin d(3) is a negative endocrine regulator of the renin-angiotensin system. *The Journal of clinical investigation*, 110(2):229–238, Jul 2002.
- [83] S. Pilz, A. Tomaschitz, E. Ritz, and T. R. Pieber. Vitamin d status and arterial hypertension: a systematic review. *Nature reviews.Cardiology*, 6(10):621–630, Oct 2009.
- [84] K. L. Morris and M. B. Zemel. 1,25-dihydroxyvitamin d3 modulation of adipocyte glucocorticoid function. *Obesity research*, 13(4):670–677, Apr 2005.
- [85] M. H. de Borst, G. Navis, R. A. de Boer, S. Huitema, L. M. Vis, W. H. van Gilst, and H. van Goor. Specific map-kinase blockade protects against renal damage in homozygous tgr(mren2)27 rats. *Laboratory investigation; a journal of technical methods and pathology*, 83(12):1761–1770, Dec 2003.
- [86] D. J. van Veldhuisen, S. Genth-Zotz, J. Brouwer, F. Boomsma, T. Netzer, A. J. Man In 't Veld, Y. M. Pinto, K. I. Lie, and H. J. Crijns. High- versus low-dose ace inhibition in chronic heart failure: a double-blind, placebo-controlled study of imidapril. *Journal of the American College of Cardiology*, 32(7):1811–1818, Dec 1998.
- [87] J. A. Krikken, G. D. Laverman, and G. Navis. Benefits of dietary sodium restriction in the management of chronic kidney disease. *Current opinion in nephrology and hypertension*, 18(6):531–538, Nov 2009.
- [88] L. Vogt, F. Waanders, F. Boomsma, D. de Zeeuw, and G. Navis. Effects of dietary sodium and hydrochlorothiazide on the antiproteinuric efficacy of losartan. *Journal of the American Society of Nephrology : JASN*, 19(5):999–1007, May 2008.
- [89] A. B. Kramer, E. F. van der Meulen, I. Hamming, H. van Goor, and G. Navis. Effect of combining ace inhibition with aldosterone blockade on proteinuria and renal damage in experimental nephrosis. *Kidney international*, 71(5):417–424, Mar 2007.
- [90] G. Nguyen, F. Delarue, C. Burckle, L. Bouzahir, T. Giller, and J. D. Sraer. Pivotal role of the renin/prorenin receptor in angiotensin ii production and cellular responses to renin. *The Journal of clinical investigation*, 109(11):1417–1427, Jun 2002.

- [91] I. Hamming, G. Navis, M. J. Kocks, and H. van Goor. Ace inhibition has adverse renal effects during dietary sodium restriction in proteinuric and healthy rats. *The Journal of pathology*, 209(1):129–139, May 2006.
- [92] M. Mizobuchi, J. Morrissey, J. L. Finch, D. R. Martin, H. Liapis, T. Akizawa, and E. Slatopolsky. Combination therapy with an angiotensin-converting enzyme inhibitor and a vitamin d analog suppresses the progression of renal insufficiency in uremic rats. *Journal of the American Society of Nephrology : JASN*, 18(6):1796–1806, Jun 2007.
- [93] P. van Paassen, D. de Zeeuw, G. Navis, and P. E. de Jong. Renal and systemic effects of continued treatment with renin inhibitor remikiren in hypertensive patients with normal and impaired renal function. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*, 15(5):637–643, May 2000.
- [94] H. H. Parving, F. Persson, J. B. Lewis, E. J. Lewis, N. K. Hollenberg, and AVOID Study Investigators. Aliskiren combined with losartan in type 2 diabetes and nephropathy. *The New England journal of medicine*, 358(23):2433–2446, Jun 5 2008.
- [95] F. Persson, P. Rossing, H. Reinhard, T. Juhl, C. D. Stehouwer, C. Schalkwijk, A. H. Danser, F. Boomsma, E. Frandsen, and H. H. Parving. Renal effects of aliskiren compared with and in combination with irbesartan in patients with type 2 diabetes, hypertension, and albuminuria. *Diabetes care*, 32(10):1873–1879, Oct 2009.
- [96] E. Yamamoto, K. Kataoka, Y. F. Dong, T. Nakamura, M. Fukuda, Y. Tokutomi, S. Matsuba, H. Nako, N. Nakagata, T. Kaneko, H. Ogawa, and S. Kim-Mitsuyama. Aliskiren enhances the protective effects of valsartan against cardiovascular and renal injury in endothelial nitric oxide synthase-deficient mice. *Hypertension*, 54(3):633–638, Sep 2009.
- [97] W. W. Batenburg, R. J. de Bruin, J. M. van Gool, D. N. Muller, M. Bader, G. Nguyen, and A. H. Danser. Aliskiren-binding increases the half life of renin and prorenin in rat aortic vascular smooth muscle cells. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 28(6):1151–1157, Jun 2008.
- [98] Z. Zhang, Y. Zhang, G. Ning, D. K. Deb, J. Kong, and Y. C. Li. Combination therapy with a1 blocker and vitamin d analog markedly ameliorates diabetic nephropathy: blockade of compensatory renin increase. *Proceedings of the National Academy of Sciences of the United States of America*, 105(41):15896–15901, Oct 14 2008.
- [99] A. B. Kramer, G. D. Laverman, H. van Goor, and G. Navis. Inter-individual differences in anti-proteinuric response to acei in established adriamycin nephrotic rats are predicted by pretreatment renal damage. *The Journal of pathology*, 201(1):160–167, Sep 2003.
- [100] G. Remuzzi, C. Zoja, E. Gagliardini, D. Corna, M. Abbate, and A. Benigni. Combining an anti-proteinuric approach with mycophenolate mofetil fully suppresses progressive nephropathy of experimental animals. *Journal of the American Society of Nephrology : JASN*, 10(7):1542–1549, Jul 1999.
- [101] V. Panichi, M. Migliori, D. Taccola, C. Filippi, L. De Nisco, L. Giovannini, R. Palla, C. Tetta, and G. Camussi. Effects of 1,25(oh)2d3 in experimental mesangial proliferative nephritis in rats. *Kidney international*, 60(1):87–95, Jul 2001.
- [102] J. M. Lemire, J. S. Adams, R. Sakai, and S. C. Jordan. 1 alpha,25-dihydroxyvitamin d3 suppresses proliferation and immunoglobulin production by normal human peripheral blood mononuclear cells. *The Journal of clinical investigation*, 74(2):657–661, Aug 1984.
- [103] W. F. Rigby, T. Stacy, and M. W. Fanger. Inhibition of t lymphocyte mitogenesis by 1,25-dihydroxyvitamin d3 (calcitriol). *The Journal of clinical investigation*, 74(4):1451–1455, Oct 1984.
- [104] J. S. Adams and M. Hewison. Unexpected actions of vitamin d: new perspectives on the regulation of innate and adaptive immunity. *Nature clinical practice. Endocrinology & metabolism*, 4(2):80–90, Feb 2008.

- [105] D. Zehnder, M. Quinkler, K. S. Eardley, R. Bland, J. Lepenies, S. V. Hughes, N. T. Raymond, A. J. Howie, P. Cockwell, P. M. Stewart, and M. Hewison. Reduction of the vitamin d hormonal system in kidney disease is associated with increased renal inflammation. *Kidney international*, 74(10):1343–1353, Nov 2008.
- [106] M. Cohen-Lahav, A. Douvdevani, C. Chaimovitz, and S. Shany. The anti-inflammatory activity of 1,25-dihydroxyvitamin d3 in macrophages. *The Journal of steroid biochemistry and molecular biology*, 103(3-5):558–562, Mar 2007.
- [107] H. Sigmundsdottir, J. Pan, G. F. Debes, C. Alt, A. Habtezion, D. Soler, and E. C. Butcher. Dcs metabolize sunlight-induced vitamin d3 to 'program' t cell attraction to the epidermal chemokine ccl27. *Nature immunology*, 8(3):285–293, Mar 2007.
- [108] J. Fritsche, K. Mondal, A. Ehrnsperger, R. Andreessen, and M. Kreutz. Regulation of 25-hydroxyvitamin d3-1 alpha-hydroxylase and production of 1 alpha,25-dihydroxyvitamin d3 by human dendritic cells. *Blood*, 102(9):3314–3316, Nov 1 2003.
- [109] E. van Etten and C. Mathieu. Immunoregulation by 1,25-dihydroxyvitamin d3: basic concepts. *The Journal of steroid biochemistry and molecular biology*, 97(1-2):93–101, Oct 2005.
- [110] J. R. Mora, M. Iwata, and U. H. von Andrian. Vitamin effects on the immune system: vitamins a and d take centre stage. *Nature reviews.Immunology*, 8(9):685–698, Sep 2008.
- [111] G. Penna and L. Adorini. 1 alpha,25-dihydroxyvitamin d3 inhibits differentiation, maturation, activation, and survival of dendritic cells leading to impaired alloreactive t cell activation. *Journal of immunology (Baltimore, Md.: 1950)*, 164(5):2405–2411, Mar 1 2000.
- [112] L. Adorini and G. Penna. Dendritic cell tolerogenicity: a key mechanism in immunomodulation by vitamin d receptor agonists. *Human immunology*, 70(5):345–352, May 2009.
- [113] M. Ghoreishi, P. Bach, J. Obst, M. Komba, J. C. Fleet, and J. P. Dutz. Expansion of antigen-specific regulatory t cells with the topical vitamin d analog calcipotriol. *Journal of immunology (Baltimore, Md.: 1950)*, 182(10):6071–6078, May 15 2009.
- [114] C. A. Redaelli, M. Wagner, D. Gunter-Duwe, Y. H. Tian, P. F. Stahel, L. Mazzucchelli, R. A. Schmid, and M. K. Schilling. 1alpha,25-dihydroxyvitamin d3 shows strong and additive immunomodulatory effects with cyclosporine a in rat renal allotransplants. *Kidney international*, 61(1):288–296, Jan 2002.
- [115] M. Courbebaisse, E. Thervet, J. C. Souberbielle, J. Zuber, D. Eladari, F. Martinez, M. F. Mamzer-Bruneel, P. Urena, C. Legendre, G. Friedlander, and D. Prie. Effects of vitamin d supplementation on the calcium-phosphate balance in renal transplant patients. *Kidney international*, 75(6):646–651, Mar 2009.
- [116] F. Heymann, C. Meyer-Schwesinger, E. E. Hamilton-Williams, L. Hammerich, U. Panzer, S. Kaden, S. E. Quaggin, J. Floege, H. J. Grone, and C. Kurts. Kidney dendritic cell activation is required for progression of renal disease in a mouse model of glomerular injury. *The Journal of clinical investigation*, 119(5):1286–1297, May 2009.
- [117] U. Thiem, G. Heinze, R. Segel, T. Perkmann, F. Kainberger, F. Muhlbacher, W. Horl, and K. Borchardt. Vita-d: cholecalciferol substitution in vitamin d deficient kidney transplant recipients: a randomized, placebo-controlled study to evaluate the post-transplant outcome. *Trials*, 10:36, May 29 2009.

